


# Aquatic Subregions of Missouri



The Grand River is one of the larger rivers within the Central Plains Aquatic Subregion.



**Legend**

- Central Plains
- Ozarks
- Mississippi Alluvial Basin



Huzzah and Courtois Creeks and their surrounding landscape are fairly typical of the Missouri Ozarks.

**Objective**

Identify groups of major drainages that drain regions with similar physiographic character and contain relatively similar assemblages in terms of ecological composition (e.g., life history strategies and physiological tolerances).

**General Description**

Aquatic Subregions are physiographic or ecoregional substrata of Regions and thus account for differences in the ecological composition of riverine assemblages resulting from geographic variation in ecosystem structure and function. The three Aquatic Subregions that cover Missouri (i.e., Central Plains, Ozarks, and Mississippi Alluvial Basin) largely correspond with the three major aquatic faunal regions of Missouri described by Pflieger (1989). Pflieger (1989) used a species distributional limit analysis and multivariate analyses of fish community data to empirically define these three major faunal regions. We slightly modified the boundaries of Pflieger's faunal regions to ensure that the boundaries between Subregions followed major drainage divides in order to account for drainage-specific evolutionary histories in succeeding levels of the hierarchy. Subsequent studies examining macroinvertebrate assemblages have provided additional empirical evidence that these Subregions are necessary strata to account for biophysical variation in Missouri's riverine ecosystems (Pflieger 1996; Rabeni et al. 1997; Rabeni and Doisy 2000). Each Subregion contains streams with relatively distinct structural features, functional processes, and aquatic assemblages in terms of both taxonomic and ecological composition.

**Principle Investigators**

Scott P. Sowa, Missouri Resource Assessment Partnership (MoRAP), UMC, Columbia, Missouri  
 Gust M. Annis, MoRAP, UMC, Columbia, Missouri  
 David D. Diamond, MoRAP, UMC, Columbia, Missouri  
 Michael E. Morey, MoRAP, UMC, Columbia, Missouri

**References**

- Pflieger, W. L. 1989. *Aquatic community classification system for Missouri*. Jefferson City, MO: Missouri Department of Conservation, Aquatic Series No. 19.
- Pflieger, W. L. 1996. *The Crayfishes of Missouri*. Missouri Department of Conservation, Jefferson City, MO.
- Rabeni, C. F., R. J. Sarver, N. Wang, G. S. Wallace, M. Weiland, and J. T. Peterson. 1997a. *Development of Regionally Based Biological Criteria for Streams of Missouri*. A report to the Missouri Department of Natural Resources from the Missouri Cooperative Fish and Wildlife Research Unit, University of Missouri, Columbia, MO.
- Rabeni, C. F., M. A. Smale, and E. B. Nelson. 1997b. Water quality and riparian conditions in the upper Niangua River Basin, 1991-1995; Effects upon fish and invertebrate communities. Final report submitted to the Missouri Department of Natural Resources, Division of Environmental Quality, Jefferson City, MO.
- Rabeni, C. F. and K. E. Doisy. 2000. The correspondence of stream benthic invertebrate communities to regional classification schemes in Missouri. *Journal of the North American Benthological Society*.19: 419-428.

3




Much of the Mississippi Alluvial Basin Aquatic Subregion originally consisted of vast swamps.

Annis G. M., S. P. Sowa, M. E. Morey, and D. D. Diamond. 2005. Aquatic Subregions of Missouri. MoRAP Map Series MS-2005-005.

Cartographer: Gust M. Annis

Photos compliment of the Missouri Department of Conservation: Copyright © Missouri Conservation Commission



Missouri Resource Assessment Partnership (MoRAP), University of Missouri, 4200 New Haven Road, Columbia, MO 65201